



FOR PHARMACY USE ONLY.
NOT FOR RETAIL SALE.

NDC 0904-5125-59

SudoGest™

NASAL DECONGESTANT 60 mg

Pseudoephedrine Hydrochloride 60 mg
Relieves Nasal and Sinus Congestion due to Colds or Hay Fever
Without Drowsiness

100 TABLETS



FOR PHARMACY USE ONLY.
NOT FOR RETAIL SALE.

NDC 0904-5125-59

SudoGest™

NASAL DECONGESTANT 60 mg

Pseudoephedrine Hydrochloride 60 mg
Relieves Nasal and Sinus Congestion due to Colds or Hay Fever
Without Drowsiness

100 TABLETS



SudoGest™

NASAL DECONGESTANT 60 mg

100 TABLETS

Distributed by MAJOR® PHARMACEUTICALS
31778 Enterprise Drive
Livonia, MI 48150 USA Rev 02/13 M-17
50844 REV07/2N11312

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING

B-1212-113-12
REV07/2N11312

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and older: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.
- children under 12 years of age: do not use

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see and flap for expiration date and lot number

Inactive ingredients corn starch, hydroxymethylcellulose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silicon dioxide, stearic acid

Questions or comments?

(800) 616-2471

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient
(in each tablet)
Pseudoephedrine HCl 60 mg decongestant

Purpose
Nasal

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves sinus congestion and pressure

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI). Certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland

When using this product do not exceed recommended dose.