

Drug Facts (continued)

Directions

- take every 4 to 6 hours ■ do not take more than 6 doses in 24 hours
- use only with enclosed dosing cup. Do not use with any other device
- find right dose on chart below ■ mL = milliliter

adults and children 12 years and over	10-20mL (25 mg to 50 mg)
children 6 to under 12 years	5-10mL (12.5 mg to 25 mg)
children 2 to 5 years of age	do not use unless directed by a doctor
children under 2 years of age	do not use

Other information

- each (5 mL) contains: **sodium 7 mg**
- **TAMPER-EVIDENT:** Do not use this product if inner foil seal over the mouth of the bottle is cut, torn, broken or missing
- store at room temperature 20°-25°C (68°-77°F)

Inactive ingredients artificial cherry flavor, citric acid, D&C Red #33, FD&C Red #40, glycerin, polysorbate 20, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol solution

Questions or comments? (800) 616-2471

Distributed by:

MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152

M-64 Rev. 10/18

Re-order No. 235192



* This product is not distributed or manufactured by Pfizer Consumer HealthCare, Pfizer, Inc. distributor of Benadryl® Allergy Liquid. Benadryl® is a registered trademark of Warner-Lambert Company L.L.C.

NDC 0904-1228-00
TAMPER-EVIDENT

MAJOR [®] **Banophen** [™]
ALLERGY

Antihistamine
12.5 mg Diphenhydramine HCl USP
ALCOHOL FREE
FOR THE TEMPORARY
RELIEF OF:
Upper Respiratory Allergies
Hay Fever
The Common Cold

Compare to the active ingredient
of Benadryl® Allergy Liquid*

4 FL. OZ. (118 mL)

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
Diphenhydramine HCl USP 12.5 mg Antihistamine

Uses temporarily relieves these symptoms of hay fever
or other upper respiratory allergies ■ runny nose ■ sneezing
■ itchy, watery eyes ■ itching of the nose or throat

Warnings

Do not use ■ to make a child sleepy ■ with any other product
containing diphenhydramine, including one applied topically.

Ask a doctor before use if you have

- glaucoma ■ a sodium restricted diet
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives
or tranquilizers.

When using this product

 ■ marked drowsiness may occur

- avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may
increase drowsiness ■ use caution when driving a motor vehicle or
operating machinery ■ excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use.
Keep out of the reach of children. In case of overdose, get medical help or contact
a Poison Control Center right away (1-800-222-1222).

PEEL HERE ▼